

TITLE - CINNARIZINE + DOMPERIDONE / STUGIL MEDICATION PATIENT INFORMATION IN ENGLISH

Disclaimer : Document presented by www.911GlobalMeds.com



Express Medicine Shipments, Globally!

FOR PUBLIC INTEREST & INFORMATION ONLY.
NO BRAND OR GENERIC MEDICINE IS BEING PROMOTED
FOR SALES FROM THE CONTENT OF THIS DOCUMENT.

Source : Lfylink.com

[Home \(/home/\)](#) > [Medicine \(/medicine-search/\)](#) > [Stugil Tablet](#)

Stugil Tablet

Manufacturer :

Janssen Pharmaceuticals

Composition :

Cinnarizine, Domperidone

Prescription :

Not Required

What are the interactions for Stugil Tablet?

At whatever point you take more than one medicine, or mix it with specific foods or beverages, you're in danger of a drug interaction.

Interaction with Disease

Kidney Disease Moderate

Stugil Tablet ought to be utilized with alert in the event that you are experiencing impeded kidney work.

Reasonable portion modification and security checking is exhorted, particularly if the debilitation is serious.

Liver Disease Moderate

Stugil Tablet ought to be utilized with alert in the event that you are experiencing impeded liver capacity.

Reasonable portion modification and wellbeing observing is exhorted, particularly if the weakness is gentle to direct. For extreme disability, utilization of this medicine isn't suggested.

Intestinal Disorders Major

Stugil Tablet ought to be utilized with extraordinary alert in individuals having genuine intestinal problems like inside dying, an obstacle or aperture of the stomach and digestive tract. Utilization of a substitute medicine is prompted in such cases.

Interaction with Alcohol

Liquor

Interaction with liquor is obscure. It is fitting to counsel your specialist before utilization.

Interaction with Lab Test

Lab

Data not accessible.

Interaction with Food

Nourishment

Data not accessible.

Interaction with Medicine

Tramadol Major

Report the use of Tramadol or some other medicine having a place with Opioid class to the specialist. Reasonable portion modification and security observing is exhorted in these cases.

Ketoconazole Major

Report the utilization of Ketoconazole or some other enemy of parasitic medicine to the specialist. Co-organization of these medicines isn't prescribed as the danger of side effects are high. Try not to cease the utilization of any drug without counseling your specialist.

Erythromycin Major

Report the utilization of Erythromycin to the specialist. Co-organization of these medicines isn't suggested as the danger of side effects are exceptionally high. Try not to stop the use of any prescription without counseling your specialist.

Bromocriptine Moderate

Report the utilization of both of the drug to the specialist. You may require a portion alteration if these medicines are to be directed together. Try not to cease the utilization of any medicine without counseling your specialist.

Cisapride Major

Report the utilization of both of the medicine to the specialist. Co-organization of these medicines isn't prescribed and utilization of an option is recommended. Try not to cease the utilization of any medicine without counseling your specialist.

Where is the Stugil Tablet approved?

India

Japan

What are the dosage instructions?

Are there any missed dose instructions?

Skip the missed dose and resume with the next scheduled dose at the right time. The dose should not get doubled in an attempt to make up for the missed dose.

Are there any overdose instructions?

Contact your doctor if an overdose is suspected. Symptoms of overdose may include drowsiness, agitation, and convulsions and may be more prevalent in infants and children. Immediate medical attention including gastric lavage may be required in severe cases.

What are the substitutes for Stugil Tablet?

Below is the list of medicines, which have the same composition, strength and form as Stugil Tablet, and hence can be used as its substitute.

Cinirone D Tablet

Biochem Pharmaceutical Industries

Kvert D 15Mg/20Mg Tablet

Kivi Labs Ltd

Cindom 15 Mg/20 Mg Tablet

Tas Med India Pvt Ltd

Vertidom Tablet

Geno Pharmaceuticals Ltd

Synadom 15 Mg/20 Mg Tablet

Bsp Pharmaceuticals

CINERAX-D 15MG/20MG TABLET

Rax Health Care Pvt Ltd

Spinil D 15 Mg/20 Mg Tablet

Alchemist Life Science Ltd

Rhizand 15 Mg/20 Mg Tablet

Cure Quick Pharmaceuticals

Domax Cz Tablet

Mother Remedies

Domfast Cz 15 Mg/20 Mg Tablet

Swiss Pharma Pvt Ltd

AUGZIN 15MG/20MG TABLET

Auriga Labs

Qvert D 15 Mg/20 Mg Tablet

Pegasus Farmaco India Pvt Ltd

Trudom 15 Mg/20 Mg Tablet

Genesis Biotech Inc

Novadom C Tablet

Pacific Drugs & Chemicals

Vertirid D 15 Mg/20 Mg Tablet

RKG Pharma

CINRON 15MG/20MG TABLET

Man Serve Pharma

Stedidom 15 Mg/20 Mg Tablet

Morepen Laboratories Ltd

Cinn D 15 Mg/20 Mg Tablet

Curagen Biological Pharmaceuticals

Domitin C 15 Mg/20 Mg Capsule

Navil Laboratories Pvt Ltd

Domgil 15 Mg/20 Mg Tablet

Kinedex Healthcare Pvt Ltd

DOMLEY-C 15MG/20MG TABLET

Zodley Pharmaceuticals Pvt Ltd

Cinidom 15 Mg/20 Mg Tablet

Eskag Pharma Pvt Ltd

Domcin 15 Mg/20 Mg Tablet

Intel Pharmaceuticals

Cinadew 15Mg/20Mg Tablet

Dew Drops Lab

Key highlights of Stugil Tablet

How long is the duration of effect?

The effect of this medicine lasts up to 6 hours.

What is the onset of action?

The peak effect of this medicine can be observed within 30 to 60 minutes of administration.

Are there any pregnancy warnings?

This medicine is not recommended for use in pregnant women unless the potential benefits outweigh the risks involved. The effect of this medicine on the fetus is not clearly established and hence you are advised to consult a doctor before taking this medicine.

Is it habit forming?

No habit forming tendencies were reported.

Are there any breast-feeding warnings?

This medicine is not recommended for use by breastfeeding mothers as there is a risk of side effects on the infant. You are advised to consult your doctors and consider the potential benefit and risks before consuming this medication.

What are the side effects of Stugil Tablet?

In addition to its intended effect, Stugil Tablet may cause some unwanted effects too. In such cases, you must seek medical attention immediately. This is not an exhaustive list of side effects. Please inform your doctor if you experience any adverse reaction to the medication.

Swelling of face, lips, eyelids, tongue, hands and feet Major Less Common

Difficulty in breathing Major Common

Skin rash Moderate Common

Convulsions Moderate Common

Heart rhythm disorders Major Rare

Disrupted menstrual cycle Moderate Rare

Breast pain and tenderness Moderate Less Common

Dry mouth Moderate Common

Loss of libido Moderate Less Common

Breast like growth in men Moderate Rare

What are the contraindications of Stugil Tablet?

Allergy

Stugil Tablet is not recommended for use in people having a known history of allergy to it.

Tumor of Pituitary gland

Stugil Tablet is not recommended for use in people having tumor of the pituitary gland.

Heart diseases

Stugil Tablet is not recommended for use in people having an active heart disease like Congestive Heart Failure.

When is Stugil Tablet prescribed?

Nausea and vomiting

Stugil Tablet is used to treat or prevent nausea and vomiting which may be caused due to chemotherapy or digestive system disorder.

Gastric motility disorders

Stugil Tablet is also used in the treatment of gastric motility disorders which may include delayed or hastened emptying of stomach.

About Stugil Tablet

Stugil Tablet, a piece of the dopamine opponents gathering of drugs, keeps the propensity of spewing and queasiness in individuals experiencing gastrointestinal problems and furthermore the individuals who take certain medicines for Parkinsons malady. The medicine fixes the muscles close to the passage of the stomach and loosens up the ones present at the exit. This activity accelerates the death of the nourishment that you consumption - from

your stomach to your digestive tract, lessening the sentiment of queasiness and disorder, additionally preventing spewing. It can likewise square, or diminish incitement in the regurgitating focus of your mind, which will in the end decrease the sentiment of queasiness and heaving.

Stugil Tablet is a piece of the drug assemble known as dopamine opponents. It very well may be utilized for treating the moderate section of sustenance in your gastrointestinal tract, for the most part connected with gastritis or diabetes. For individuals experiencing this condition, Stugil Tablet can treat the indications of spewing, sickness, swelling and feeling full. Aside from that, it can likewise forestall spewing and queasiness that is related with Parkinsons sickness. The prescription works by discharging your stomach rapidly, decreasing queasiness. It additionally lessens or squares incitement in the piece of the cerebrum called the heaving focus. The nerve messages originating from your gut will be stifled and the sentiment of queasiness and regurgitating is counteracted. It is accessible in tablet or suspension structure and must be taken orally.

The typical portion of Stugil Tablet is 10 mg, which is commonly taken 15-30 minutes preceding your suppers. The greatest portion that is prescribed for grown-ups is 30 mg. The portion that would suit your body relies upon your body weight, different medicines that you are right now taking and your wellbeing condition. Guarantee that you take Stugil Tablet just in that sum as taught by the specialist. In the event that you miss a portion, you can take it later. Nonetheless, don't take two portions together to compensate for the past one that has been missed.

Overdose can cause bewilderment, dizziness, losing muscle or parity control or trouble in talking.

You are prompted not to take Stugil Tablet on the off chance that you have the accompanying conditions:

A hypersensitivity to Stugil Tablet or some other element of this medicine

Draining problems or blockage in your stomach or in the digestive tract

Tumor in the pituitary organ

Heart sickness

Inaccurate dimensions of magnesium, potassium or calcium in the blood

Extreme/moderate liver debilitation

The conceivable side effects of taking Stugil Tablet can be looseness of the bowels, cerebral pain, headache, dryness of the mouth or bosom torment. These side effects are not extremely normal and leave in a couple of days.

If not, you can counsel your specialist. Aside from these, you should look for restorative consideration immediately and quit taking Stugil Tablet in the event that you experience any of the accompanying side effects:

Inconsistency in heart beat

A sentiment of unsteadiness or faintness

Trouble in breathing, swelling of the throat or the face

Menstrual inconsistencies

Release of bosom milk from the areola

Swelling of bosoms in guys

You do not have any consultation yet.

[Ask a doctor \(/askquestion/\)](/askquestion/)

Did you find this helpful?

Your feedback will help us improve the product

Yes

No

0 people found it helpful

(<https://www.facebook.com/lyfLink.in>)

(<https://twitter.com/lyfLink/>)

(<https://plus.google.com/112483408314968322009>)

Information on this page was last updated on Jun 25, 2019

Report an error

Looking For a Doctor?

Book an appointment with the top doctors near you.

[Book Now \(/search/doctor/\)](/search/doctor/)

Disclaimer

We've made all possible efforts to ensure that the information provided here is accurate, up-to-date and complete, however, it should not be treated as a substitute for professional medical advice, diagnosis or treatment. LyfLink only provides reference source for common information on medicines and does not guarantee its accuracy or exhaustiveness. The absence of a warning for any drug or combination thereof, should not be assumed to indicate that the drug or combination is safe, effective, or appropriate for any given patient. LyfLink does not assume any responsibility for any aspect of healthcare administered with the aid of information provided above. If you have any doubts about your medication then we strongly recommend that you consult with your doctor, nurse or healthcare provider. See detailed T&C [here \(/medicine-terms-and-conditions/\)](/medicine-terms-and-conditions/).